arXiv:1410.7205v1 [math.CA] 5 Oct 2014

STRONG CONVERGENCE OF TWO–DIMENSIONAL WALSH-FOURIER SERIES

G. TEPHNADZE

Abstract. We prove that certain mean of the quadratical partial sums of the two-dimensional Walsh-Fourier series are uniformly bounded operators from the Hardy space H_p to the space L_p for $0 < p < 1$.

2010 Mathematics Subject Classification. 42C10.

Key words and phrases: Walsh system, Strong convergence, martingale Hardy space.

1. INTRODUCTION

It is known [7, p. 125] that the Walsh-Paley system is not a Schauder basis in $L_1(G)$. Moreover, (see [8]) there exists a function in the dyadic Hardy space $H_1(G)$, the partial sums of which are not bounded in $L_1(G)$. However, in Simon [9] the following strong convergence result was obtained for all $f \in H_1$:

$$\lim_{n\to\infty} \frac{1}{\log n} \sum_{k=1}^{n} \frac{\|S_k f - f\|_1}{k} = 0,$$

where $S_k f$ denotes the $k-$th partial sum of the Walsh-Fourier series of f (For the trigonometric analogue see Smith [11], for the Vilenkin system see Gát [1]).

Simon [10] proved that there is an absolute constant c_p, depends only p, such that

$$\sum_{k=1}^{\infty} \frac{\|S_k f\|_p^p}{k^{2-p}} \leq c_p \|f\|_{H_p}^p, \tag{1}$$

for all $f \in H_p$, where $0 < p < 1$.

The author [13] proved that sequence $\{1/k^{2-p}\}_{k=1}^{\infty}$ in inequality (1) is important.

For the two-dimensional Walsh-Fourier series Weisz [16] generalized the result of Simon and proved that if $\alpha \geq 0$ and $f \in H_p(G \times G)$, then

$$\sup_{n,m\geq 2} \left(\frac{1}{\log n \log m}\right)^{[p]} \sum_{2^{-\alpha}\leq k/l\leq 2^{\alpha},\ (k,l)\leq(n,m)} \frac{\|S_{k,l} f\|_p^p}{(kl)^{2-p}} \leq c \|f\|_{H_p}^p,$$

where $0 < p < 1$ and $[p]$ denotes the integer part of p.

Goginava and Gogoladze [5] proved that the following result is true:

Theorem G.Let $f \in H_1(G \times G)$. Then there exists absolute constant c, such that

$$\sum_{n=1}^{\infty} \frac{\|S_{n,n} f\|_1}{n \log^2 n} \leq c \|f\|_{H_1}.$$

For two-dimensional trigonometric system analogical theorem was proved in [6].

Convergence of quadratical partial sums of two-dimensional Walsh-Fourier series was investigated in details by Weisz [15], Goginava [4], Gát, Goginava, Nagy [2], Gát, Goginava, Tkebuchava [3].

The main aim of this paper is to prove (see Theorem 1) that

$$\sum_{n=1}^{\infty}\frac{\|S_{n,n}f\|_p^p}{n^{3-2p}}\leq c_p\|f\|_{H_p}^p, \tag{2}$$

for all $f\in H_p(G\times G)$, where $0<p<1$. We also proved that sequence $\{1/n^{3-2p}\}_{n=1}^{\infty}$ in inequality 2 is important (see Theorem 2).

2. DEFINITIONS AND NOTATIONS

Let $\mathbf{P}$ denote the set of positive integers, $\mathbf{N}:=\mathbf{P}\cup\{0\}$. Denote by Z_2 the discrete cyclic group of order 2, that is $Z_2=\{0,1\}$, where the group operation is the modulo 2 addition and every subset is open. The Haar measure on Z_2 is given such that the measure of a singleton is $1/2$. Let G be the complete direct product of the countable infinite copies of the compact groups Z_2. The elements of G are of the form $x=(x_0,x_1,...,x_k,...)$ with $x_k\in\{0,1\}\,(k\in\mathbf{N})$. The group operation on G is the coordinate-wise addition, the measure (denote by μ) and the topology are the product measure and topology. The compact Abelian group G is called the Walsh group. A base for the neighborhoods of G can be given in the following way:

$$\begin{aligned}I_0(x) &:= G,\ \ I_n(x):=I_n(x_0,...,x_{n-1})\\ &:=\{y\in G:\ y=(x_0,...,x_{n-1},y_n,y_{n+1},...)\},\\ &\qquad (x\in G,n\in\mathbf{N}).\end{aligned}$$

These sets are called the dyadic intervals. Let $0=(0:i\in\mathbf{N})\in G$ denote the null element of G, $I_n:=I_n(0)\ (n\in\mathbf{N})$. Set $e_n:=(0,...,0,1,0,...)\in G$ the n th coordinate of which is 1 and the rest are zeros $(n\in\mathbf{N})$. Let $\overline{I}_n:=G\backslash I_n$.

If $n\in\mathbf{N}$, then $n=\sum_{i=0}^{\infty}n_i2^i$, where $n_i\in\{0,1\}\ \ (i\in\mathbf{N})$, i. e. n is expressed in the number system of base 2. Denote $|n|:=\max\{j\in\mathbf{N}:n_j\neq 0\}$, that is, $2^{|n|}\leq n<2^{|n|+1}$.

It is easy to show that for every odd number $n_0=1$ and we can write $n=1+\sum_{i=1}^{|n|}n_j2^i$, where $n_j\in\{0,1\}$, $(j\in\mathbf{P})$.

For $k\in\mathbf{N}$ and $x\in G$ let as denote by

$$r_k(x):=(-1)^{x_k}\qquad (x\in G,k\in\mathbf{N})$$

the k-th Rademacher function.

The Walsh-Paley system is defined as the sequence of Walsh-Paley functions:

$$w_n(x):=\prod_{k=0}^{\infty}(r_k(x))^{n_k}=r_{|n|}(x)(-1)^{\sum_{k=0}^{|n|-1}n_kx_k}\qquad (x\in G,n\in\mathbf{P}).$$

The Walsh-Dirichlet kernel is defined by

$$D_n(x)=\sum_{k=0}^{n-1}w_k(x).$$

Recall that (see [8, p. 7])

$$D_{2^n}(x) = \begin{cases} 2^n, & x \in I_n \\ 0, & x \in \overline{I}_n \end{cases}, \tag{3}$$

Furthermore, the following representation holds for the D_n's. Let $n \in \mathbf{N}$ and $n = \sum_{i=0}^{\infty} n_i 2^i$. Then

$$D_n(x) = w_n(x) \sum_{j=0}^{\infty} n_j w_{2^j}(x) D_{2^j}(x). \tag{4}$$

The rectangular partial sums of the 2-dimensional Walsh-Fourier series of function $f \in L_2(G \times G)$ are defined as follows:

$$S_{M,N} f(x,y) := \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} \widehat{f}(i,j) w_i(x) w_j(y),$$

where the numbers

$$\widehat{f}(i,j) = \int_{G \times G} f(x,y) w_i(x) w_j(y) d\mu(x,y)$$

is said to be the $(i,j)-$th Walsh-Fourier coefficient of the function f.

Denote

$$S_M^{(1)} f(x,y) := \int_G f(s,y) D_M(x+s) d\mu(s)$$

and

$$S_N^{(2)} f(x,y) := \int_G f(x,t) D_N(y+t) d\mu(t).$$

The norm (or quasinorm) of the space $L_p(G \times G)$ is defined by

$$\|f\|_p := \left(\int_{G \times G} |f|^p d\mu \right)^{1/p}, \qquad (0 < p < \infty).$$

The space $weak - L_p(G \times G)$ consists of all measurable functions f for which

$$\|f\|_{weak-L_p(G\times G)} := \sup_{\lambda>0} \lambda \mu(f > \lambda)^{1/p} < +\infty.$$

The $\sigma-$algebra generated by the dyadic 2-dimensional $I_n(x) \times I_n(y)$ square of measure $2^{-n} \times 2^{-n}$ will be denoted by $F_{n,n} (n \in \mathbf{N})$. Denote by $f = (f_{n,n}\ n \in \mathbf{N})$ one-parameter martingale with respect to $F_{n,n} (n \in \mathbf{N})$.(for details see e.g. [14]).

The expectation operator and the conditional expectation operator relative to the $F_{n,n} (n \in \mathbf{N})$ are denoted by E and $E_{n,n}$, respectively.

The maximal function of a martingale f is defined by

$$f^* = \sup_{n \in \mathbf{N}} |f_{n,n}|.$$

Let $f\in L_1\left(G\times G\right)$. Then the dyadic maximal function is given by

$$f^{*}\left(x,y\right)=\sup_{n\in\mathbf{N}}\frac{1}{\mu\left(I_n(x)\times I_n(y)\right)}\left|\int\limits_{I_n(x)\times I_n(y)}f\left(s,t\right)d\mu\left(s,t\right)\right|,$$

$$(x,y)\in G\times G.$$

The dyadic Hardy space $H_p(G\times G)$ $(0<p<\infty)$ consists of all functions for which

$$\left\|f\right\|_{H_p}:=\left\|f^{*}\right\|_{p}<\infty.$$

If $f\in L_1\left(G\times G\right)$, then it is easy to show that the sequence $\left(S_{2^n,2^n}\left(f\right):n\in\mathbf{N}\right)$ is a martingale. If $f=\left(f_{n,n},n\in\mathbf{N}\right)$ is a martingale, then the Walsh-Fourier coefficients must be defined in a slightly different manner:

$$\widehat{f}\left(i,j\right):=\lim_{k\to\infty}\int_{G}f_{k,k}\left(x,y\right)w_i\left(x\right)w_j\left(y\right)d\mu\left(x,y\right).$$

It is known [12] that that Fourier coefficients of $f\in H_p\left(G\times G\right)$ are not bounded when $0<p<1$.

The Walsh-Fourier coefficients of $f\in L_1\left(G\times G\right)$ are the same as those of the martingale $\left(S_{2^n,2^n}f:n\in\mathbf{N}\right)$ obtained from f .

A bounded measurable function a is a p-atom, if there exists a dyadic 2-dimensional cube $I\times I$, such that

$$\begin{cases} a) & \int_{I\times I}ad\mu=0,\\ b) & \left\|a\right\|_{\infty}\leq\mu(I\times I)^{-1/p},\\ c) & \operatorname{supp}\left(a\right)\subset I\times I. \end{cases}$$

3. FORMULATION OF MAIN RESULTS

Theorem 1. *Let $0<p<1$ and $f\in H_p\left(G\times G\right)$. Then*

$$\sum_{n=1}^{\infty}\frac{\left\|S_{n,n}f\right\|_{p}^{p}}{n^{3-2p}}\leq c_p\left\|f\right\|_{H_p}^{p}.$$

Theorem 2. *Let $0<p<1$ and $\Phi:\mathbf{N}\to[1,\infty)$ is any nondecreasing function, satisfying the condition $\lim\limits_{n\to\infty}\Phi\left(n\right)=+\infty$. Then there exists a martingale $f\in H_p\left(G\times G\right)$ such that*

$$\sum_{n=1}^{\infty}\frac{\left\|S_{n,n}f\right\|_{weak-L_p}^{p}\Phi\left(n\right)}{n^{3-2p}}=\infty.$$

4. AUXILIARY PROPOSITIONS

Lemma 1. [14] *A martingale $f\in L_p\left(G\times G\right)$ is in $H_p\left(G\times G\right)\left(0<p\leq1\right)$ if and only if there exist a sequence $\left(a_k,k\in\mathbf{N}\right)$ of p-atoms and a sequence $\left(\mu_k,k\in\mathbf{N}\right)$ of a real numbers such that*

$$\sum_{k=0}^{\infty}\mu_kE_{n,n}a_k=f_{n,n}\tag{5}$$

and

$$\sum_{k=0}^{\infty} |\mu_k|^p < \infty,$$

Moreover, $\|f\|_{H_p} \backsim \inf \left(\sum_{k=0}^{\infty} |\mu_k|^p\right)^{1/p}$, where the infimum is taken over all decomposition of f of the form (5).

5. PROOF OF THE THEOREMS

Proof of Theorem 1. If we apply Lemma 1 we only have to prove that

$$\sum_{n=1}^{\infty} \frac{\|S_{n,n}a\|_p^p}{n^{3-2p}} \leq c_p < \infty, \tag{6}$$

for every p atom a.

Let a be an arbitrary p-atom with support $I_N(z') \times I_N(z'')$ and $\mu(I_N) = \mu(I_N) = 2^{-N}$. We can suppose that $z' = z'' = 0$.

Let $(x, y) \in \overline{I}_N \times \overline{I}_N$. In this case $D_{2^i}(x+s) 1_{I_N}(s) = 0$ and $D_{2^i}(y+t) 1_{I_N}(t) = 0$ for $i \geq N$. Recall that $w_{2^j}(x+t) = w_{2^j}(x)$ for $t \in I_N$ and $j < N$. Consequently, from (4) we obtain

$$\begin{aligned}
& S_{n,n}a(x,y) \\
= & \int_{G\times G} a(s,t) D_n(x+s) D_n(y+t) d\mu(s,t) \\
= & \int_{I_N\times I_N} a(s,t) D_n(x+s) D_n(y+t) d\mu(s,t) \\
= & \int_{I_N\times I_N} a(s,t) w_n(x+s+y+t) \sum_{i=0}^{N-1} n_i w_{2^i}(x+s) D_{2^i}(x+s) \\
& \times \sum_{j=0}^{N-1} n_j w_{2^j}(y+t) D_{2^j}(y+t) d\mu(s,t) \\
= & \ w_n(x) \sum_{i=0}^{N-1} n_i w_{2^i}(x) D_{2^i}(x) w_n(y) \sum_{j=0}^{N-1} n_j w_{2^j}(y) D_{2^j}(y) \\
& \times \int_{I_N\times I_N} a(s,t) w_n(s+t) d\mu(s,t)
\end{aligned}$$

$$
\begin{aligned}
&= w_n(x+y)\sum_{i=0}^{N-1} n_i w_{2^i}(x) D_{2^i}(x)\sum_{j=0}^{N-1} n_j w_{2^j}(y) D_{2^j}(y)\\
&\quad\times\int_{I_N}\left(\int_{I_N} a(t+\tau,t)\,d\mu(t)\right) w_n(\tau)\,d\mu(\tau)\\
&= w_n(x+y)\sum_{i=0}^{N-1} n_i w_{2^i}(x) D_{2^i}(x)\sum_{j=0}^{N-1} n_j w_{2^j}(y) D_{2^j}(y)\int_{I_N}\Phi(\tau)\, w_n(\tau)\,d\mu(\tau)\\
&= w_n(x+y)\sum_{i=0}^{N-1} n_i w_{2^i}(x) D_{2^i}(x)\sum_{j=0}^{N-1} n_j w_{2^j}(y) D_{2^j}(y)\,\widehat{\Phi}(n),
\end{aligned}
$$

where

$$\Phi(\tau)=\int_{I_N} a(t+\tau,t)\,d\mu(t).$$

Let $x\in I_s\backslash I_{s+1}$. Using (3) we get

$$\sum_{i=0}^{N-1} D_{2^i}(x)\le c2^s.$$

Since

$$\overline{I}_N=\bigcup_{s=0}^{N-1} I_s\backslash I_{s+1}$$

we obtain

$$
\begin{aligned}
&\int_{\overline{I}_N}\left(\sum_{i=0}^{N-1} D_{2^i}(x)\right)^p d\mu(x)\\
&\le c_p\sum_{s=0}^{N-1}\int_{I_s\backslash I_{s+1}} 2^{ps}d\mu(x)\\
&\le c_p\sum_{s=0}^{\infty}2^{(p-1)s}\\
&< c_p<\infty,\qquad 0<p<1.
\end{aligned}
\tag{7}
$$

applying (7) we can write

$$
\begin{aligned}
&\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int_{\overline{I}_N\times\overline{I}_N}|S_{n,n}a(x,y)|^p\,d\mu(x,y)\\
&\le \sum_{n=1}^{\infty}\frac{\left|\widehat{\Phi}(n)\right|^p}{n^{3-2p}}\left(\int_{\overline{I}_N}\left(\sum_{i=0}^{N-1} D_{2^i}(x)\right)^p d\mu(x)\right)^2\\
&\le c_p\sum_{n=1}^{\infty}\frac{\left|\widehat{\Phi}(n)\right|^p}{n^{3-2p}}.
\end{aligned}
$$

Let $n < 2^N$. Since $w_n(\tau) = 1$, for $\tau \in I_N$ we have

$$\begin{aligned}\widehat{\Phi}(n) &= \int_{I_N} \Phi(\tau) w_n(\tau) \, d\mu(\tau) \\ &= \int_{I_N} \left(\int_{I_N} a(t+\tau, t) \, d\mu(t) \right) w_n(\tau) \, d\mu(\tau) \\ &= \int_{I_N \times I_N} a(s,t) \, d\mu(s,t) = 0.\end{aligned}$$

Hence, we can suppose that $n \geq 2^N$. By Hölder inequality we obtain

$$\begin{aligned}&\sum_{n=1}^{\infty} \frac{\left|\widehat{\Phi}(n)\right|^p}{n^{3-2p}} \\ &\leq \left(\sum_{n=2^N}^{\infty} \left|\widehat{\Phi}(n)\right|^2 \right)^{p/2} \left(\sum_{n=2^N}^{\infty} \frac{1}{n^{(3-2p)\cdot(2/(2-p))}} \right)^{(2-p)/2} \\ &\leq \left(\frac{1}{2^{N(2(3-2p)/(2-p)-1)}} \right)^{(2-p)/2} \left(\int_G |\Phi(\tau)|^2 \, d\mu(\tau) \right)^{p/2} \\ &\leq \frac{c_p}{2^{N(4-3p)/2}} \left(\int_{I_N} \left| \int_{I_N} a(t+\tau, t) \, d\mu(t) \right|^2 d\mu(\tau) \right)^{p/2} \\ &\leq \frac{c_p}{2^{N(4-3p)/2}} \|a\|_\infty^p \frac{1}{2^{Np/2}} \frac{1}{2^{Np}} \\ &\leq \frac{c_p}{2^{N(4-3p)/2}} 2^{2N} \frac{1}{2^{3pN/2}} < c_p < \infty.\end{aligned} \tag{8}$$

Let $(x, y) \in \overline{I}_N \times I_N$. Then we have

$$\begin{aligned}&S_{n,n} a(x,y) \\ &= w_n(x) \sum_{j=0}^{N-1} n_j w_{2^j}(x) D_{2^j}(x) \\ &\quad \times \int_{G \times G} a(s,t) w_n(s) D_n(y+t) \, d\mu(s,t) \\ &= w_n(x) \sum_{j=0}^{N-1} n_j w_{2^j}(x) D_{2^j}(x) \int_G S_n^{(2)} a(s,y) w_n(s) \, d\mu(s) \\ &= w_n(x) \sum_{j=0}^{N-1} n_j w_{2^j}(x) D_{2^j}(x) \widehat{S}_n^{(2)} a(n,y).\end{aligned}$$

Using (7) we get

$$\begin{aligned}
&\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int\limits_{\overline{I}_N\times I_N}|S_{n,n}a\left(x,y\right)|^p\,d\mu\left(x,y\right)\\
\leq\ &\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int\limits_{\overline{I}_N\times I_N}\left(\sum_{j=0}^{N-1}D_{2^j}\left(x\right)\left|\widehat{S}_n^{(2)}a\left(n,y\right)\right|\right)^p d\mu\left(x,y\right)\\
\leq\ &\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int_{\overline{I}_N}\left(\sum_{i=0}^{N-1}D_{2^i}\left(x\right)\right)^p d\mu\left(x\right)\cdot\int\limits_{I_N}\left|\widehat{S}_n^{(2)}a\left(n,y\right)\right|^p d\mu\left(y\right)\\
\leq\ &\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int\limits_{I_N}\left|\widehat{S}_n^{(2)}a\left(n,y\right)\right|^p d\mu\left(y\right).
\end{aligned}$$

Let $n<2^N$. Then by the definition of the atom we have

$$\begin{aligned}
\widehat{S}_n^{(2)}a\left(n,y\right)\ &=\ \int\limits_G\left(\int\limits_G a\left(s,t\right)D_n\left(y+t\right)d\mu\left(t\right)\right)w_n\left(s\right)d\mu\left(s\right)\\
&=\ D_n\left(y\right)\int\limits_{I_N\times I_N}a\left(s,t\right)d\mu\left(s,t\right)=0.
\end{aligned}$$

Therefore, we can suppose that $n\geq 2^N$. Hence

$$\begin{aligned}
&\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int\limits_{\overline{I}_N\times I_N}|S_{n,n}a\left(x,y\right)|^p\,d\mu\left(x,y\right)\\
\leq\ &\sum_{n=2^N}^{\infty}\frac{1}{n^{3-2p}}\int\limits_{I_N}\left|\widehat{S}_n^{(2)}a\left(n,y\right)\right|^p d\mu\left(y\right)
\end{aligned}$$

Since

$$\left\|S_n^{(2)}a\left(n,y\right)\right\|_2\leq c\left\|a\right\|_2$$

from Hölder inequality we can write

$$\begin{aligned}
&\int\limits_{I_N}\left|\widehat{S}_n^{(2)}a\left(n,y\right)\right|^p d\mu\left(y\right)\\
\leq\ &\frac{c_p}{2^{N(1-p)}}\left(\int\limits_{I_N}\left|\widehat{S}_n^{(2)}a\left(n,y\right)\right|d\mu\left(y\right)\right)^p\\
=\ &\frac{c_p}{2^{N(1-p)}}\left(\int\limits_{I_N}\left|\int\limits_{I_N}S_n^{(2)}a\left(s,y\right)w_n\left(s\right)d\mu\left(s\right)\right|d\mu\left(y\right)\right)^p
\end{aligned}$$

$$
\begin{aligned}
&= \frac{c_p}{2^{N(1-p)}} \left(\int_{I_N} \left| \int_{I_N} \left(\int_{I_N} a(s,t)\, D_n(y+t)\, d\mu(t) \right) w_n(s)\, d\mu(s) \right| d\mu(y) \right)^p \\
&\le \frac{c_p}{2^{N(1-p)}} \left(\int_{I_N} \left(\int_{I_N} \left| \int_{I_N} a(s,t)\, D_n(y+t)\, d\mu(t) \right| d\mu(y) \right) d\mu(s) \right)^p \\
&\le \frac{c_p}{2^{N(1-p)}} \left(\frac{1}{2^{N/2}} \int_{I_N} \left(\int_{I_N} \left| \int_{I_N} a(s,t)\, D_n(y+t)\, d\mu(t) \right|^2 d\mu(y) \right)^{1/2} d\mu(s) \right)^p \\
&\le \frac{c_p}{2^{N(1-p)}} \left(\frac{1}{2^{N/2}} \int_{I_N} \left(\int_{I_N} |a(s,t)|^2\, d\mu(t) \right)^{1/2} d\mu(s) \right)^p \\
&\le \frac{c_p}{2^{N(1-p)}} \left(\frac{\|a\|_\infty}{2^{N/2}} \frac{1}{2^N} \frac{1}{2^{N/2}} \right)^p \le \frac{c_p}{2^{N(1-p)}} \left(\frac{2^{2N/p}}{2^{2N}} \right)^p \le c_p 2^{N(1-p)}.
\end{aligned}
$$

Consequently,

$$
\begin{aligned}
&\sum_{n=1}^{\infty} \frac{1}{n^{3-2p}} \int_{\overline{I}_N \times I_N} |S_{n,n} a(x,y)|\, d\mu(x,y) \\
&\le c_p \sum_{n=2^N}^{\infty} \frac{1}{n^{3-2p}} 2^{N(1-p)} \le \frac{c_p}{2^{N(1-p)}} \le c_p < \infty.
\end{aligned} \tag{9}
$$

Analogously, we can prove that

$$
\sum_{n=1}^{\infty} \frac{1}{n^{3-2p}} \int_{I_N \times \overline{I}_N} |S_{n,n} a(x,y)|^p\, d\mu(x,y) \le c_p < \infty. \tag{10}
$$

Let $(x,y) \in I_N \times I_N$. Then by the definition of the atom we can write

$$
\begin{aligned}
&\int_{I_N \times I_N} |S_{n,n} a(x,y)|^p\, d\mu(x,y) \\
&\le \frac{1}{2^{N(2-p)}} \left(\int_{I_N \times I_N} |S_{n,n} a(x,y)|^2\, d\mu(x,y) \right)^{p/2} \\
&\le \frac{1}{2^{N(2-p)}} \left(\int_{I_N \times I_N} |a(x,y)|^2\, d\mu(x,y) \right)^{p/2} \\
&\le \frac{\|a\|_\infty^p}{2^{N(2-p)}} \frac{1}{2^{Np}} \le c_p \frac{1}{2^{N(2-p)}} 2^{2N} \frac{1}{2^{Np}} \le c_p < \infty.
\end{aligned}
$$

It follows that

$$\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\int\limits_{I_N\times I_N}\left|S_{n,n}a\left(x,y\right)\right|d\mu\left(x,y\right) \tag{11}$$

$$\leq \quad c_p\sum_{n=1}^{\infty}\frac{1}{n^{3-2p}}\leq c_p<\infty.$$

Combining (6-11) we complete the proof of Theorem 1.

Proof of Theorem 2. Let $0<p<1$ and $\Phi\left(n\right)$ is any nondecreasing, nonnegative function, satisfying condition

$$\lim_{n\rightarrow\infty}\Phi\left(n\right)=\infty,$$

For this function $\Phi\left(n\right)$, there exists an increasing sequence of the positive integers $\{\alpha_k:\ k\geq 0\}$ such that:

$$\alpha_0\geq 2$$

and

$$\sum_{k=0}^{\infty}\frac{1}{\Phi^{p/4}\left(2^{\alpha_k}\right)}<\infty. \tag{12}$$

Let

$$f_{A,A}\left(x,y\right)=\sum_{\{k;\ \alpha_k<A\}}\lambda_k a_k,$$

where

$$\lambda_k=\frac{1}{\Phi^{1/4}\left(2^{\alpha_k}\right)}$$

and

$$a_k\left(x,y\right)=2^{\alpha_k(2/p-2)}\left(D_{2^{\alpha_k+1}}\left(x\right)-D_{2^{\alpha_k}}\left(x\right)\right)\left(D_{2^{\alpha_k+1}}\left(y\right)-D_{2^{\alpha_k}}\left(y\right)\right).$$

It is easy to show that the martingale $f=\left(f_{1,1},\ f_{2,2},...,\ f_{A,A},\ ...\right)\in H_p$.

Indeed, since

$$S_{2^A}a_k\left(x,y\right)=\begin{cases}a_k\left(x,y\right), & \alpha_k<A,\\ 0, & \alpha_k\geq A,\end{cases} \tag{13}$$

$$\begin{aligned}\mathrm{supp}(a_k) &= I_{\alpha_k},\\ \int_{I_{\alpha_k}}a_k d\mu &= 0\end{aligned}$$

and

$$\left\|a_k\right\|_{\infty}\leq 2^{\alpha_k(2/p-2)}2^{2\alpha_k}\leq 2^{2\alpha_k/p}=\left(\mathrm{supp}\ a_k\right)^{-1/p}$$

from Lemma 1 and (12) we conclude that $f\in H_p$.

It is easy to show that

$$\widehat{f}(i,j)=\begin{cases}\frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}(2^{\alpha_k})},\\ \text{if } (i,\ j)\in\{2^{\alpha_k},...,\ 2^{\alpha_k+1}-1\}\times\{2^{\alpha_k},...,\ 2^{\alpha_k+1}-1\},\ k=0,1,2...\\ 0,\\ \text{if } (\,i,\ j)\notin\bigcup\limits_{k=1}^{\infty}\{2^{\alpha_k},...,\ 2^{\alpha_k+1}-1\}\ \times\{2^{\alpha_k},...,\ 2^{\alpha_k+1}-1\}.\end{cases}\tag{14}$$

Let $2^{\alpha_k}<n<2^{\alpha_k+1}$. From (14) we have

$$\begin{aligned}&S_{n,n}f(x,y)\\ =&\sum_{i=0}^{2^{\alpha_{k-1}+1}-1}\sum_{j=0}^{2^{\alpha_{k-1}+1}-1}\widehat{f}(i,j)w_i(x)\,w_j(y)\\ &+\sum_{i=2^{\alpha_k}}^{n-1}\sum_{j=2^{\alpha_k}}^{n-1}\widehat{f}(i,j)w_i(x)\,w_j(y)\\ =&\sum_{\eta=0}^{k-1}\sum_{i=2^{\alpha_\eta}}^{2^{\alpha_\eta+1}-1}\sum_{j=2^{\alpha_\eta}}^{2^{\alpha_\eta+1}-1}\widehat{f}(i,j)w_i(x)\,w_j(y)\\ &+\sum_{i=2^{\alpha_k}}^{n-1}\sum_{j=2^{\alpha_k}}^{n-1}\widehat{f}(i,j)w_i(x)\,w_j(y)\\ =&\sum_{\eta=0}^{k-1}\sum_{i=2^{\alpha_\eta}}^{2^{\alpha_\eta+1}-1}\sum_{j=2^{\alpha_\eta}}^{2^{\alpha_\eta+1}-1}\frac{2^{\alpha_\eta(2/p-2)}}{\Phi^{1/4}(2^{\alpha_\eta})}w_i(x)\,w_j(y)\\ &+\sum_{i=2^{\alpha_k}}^{n-1}\sum_{j=2^{\alpha_k}}^{n-1}\frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}(2^{\alpha_k})}w_i(x)\,w_j(y)\\ =&\sum_{\eta=0}^{k-1}\frac{2^{\alpha_\eta(2/p-2)}}{\Phi^{1/4}(2^{\alpha_\eta})}\left(D_{2^{\alpha_\eta+1}}(x)-D_{2^{\alpha_\eta}}(x)\right)\left(D_{2^{\alpha_\eta+1}}(y)-D_{2^{\alpha_\eta}}(y)\right)\\ &+\frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}(2^{\alpha_k})}\left(D_n(x)-D_{2^{\alpha_k}}(x)\right)\left(D_n(y)-D_{2^{\alpha_k}}(y)\right)\\ =&\ I+II.\end{aligned}\tag{15}$$

Let $(x,y)\in(G\backslash I_1)\times(G\backslash I_1)$ and n is odd number. Since $n-2^{\alpha_k}$ is odd number too and

$$D_{n+2^{\alpha_k}}(x)=D_{2^{\alpha_k}}(x)+w_{2^{\alpha_k}}(x)\,D_n(x),\ \text{when}\ \ n<2^{\alpha_k},$$

from (3) and (4) we can write

$$
\begin{aligned}
(16)\qquad |II| &= \frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}\left(2^{\alpha_k}\right)}\left|w_{2^{\alpha_k}}\left(x\right)D_{n-2^{\alpha_k}}\left(x\right)w_{2^{\alpha_k}}\left(y\right)D_{n-2^{\alpha_k}}\left(y\right)\right| \\
&= \frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}\left(2^{\alpha_k}\right)}\left|w_{2^{\alpha_k}}\left(x\right)w_{n-2^{\alpha_k}}\left(x\right)D_{_1}\left(x\right)w_{2^{\alpha_k}}\left(y\right)w_{n-2^{\alpha_k}}\left(y\right)D_{_1}\left(y\right)\right| \\
&= \frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}\left(2^{\alpha_k}\right)}.
\end{aligned}
$$

Applying (3) and condition $\alpha_n \geq 2$ ($n \in \mathbf{N}$) for I we have

$$
(17)\qquad I=\sum_{\eta=0}^{k-1}\frac{2^{\alpha_k(2/p-2)}}{\Phi^{1/4}\left(2^{\alpha_\eta}\right)}\left(D_{2^{\alpha_\eta+1}}\left(x\right)-D_{2^{\alpha_\eta}}\left(x\right)\right)\left(D_{2^{\alpha_\eta+1}}\left(y\right)-D_{2^{\alpha_\eta}}\left(y\right)\right)=0.
$$

Hence

$$
\begin{aligned}
(18)\qquad &\left\|S_{n,n}f\left(x,y\right)\right\|_{weak-L_p} \\
&\geq \frac{2^{\alpha_k(2/p-2)}}{2\Phi^{1/4}\left(2^{\alpha_k}\right)}\left(\mu\left\{(x,y)\in(G\backslash I_1)\times(G\backslash I_1):\left|S_{n,n}f\left(x,y\right)\right|\geq\frac{2^{\alpha_k(2/p-2)}}{2\Phi^{1/4}\left(2^{\alpha_k}\right)}\right\}\right)^{1/p} \\
&\geq \frac{2^{\alpha_k(2/p-2)}}{2\Phi^{1/4}\left(2^{\alpha_k}\right)}\left|(G\backslash I_1)\times(G\backslash I_1)\right|\geq\frac{c_p2^{\alpha_k(2/p-2)}}{\Phi^{1/4}\left(2^{\alpha_k}\right)}.
\end{aligned}
$$

Using (18) we have

$$
\begin{aligned}
(19)\qquad &\sum_{n=1}^{2^{\alpha_k+1}-1}\frac{\left\|S_{n,n}f\right\|_{weak-L_p}^p\Phi\left(n\right)}{n^{3-2p}} \\
&\geq \sum_{n=2^{\alpha_k}+1}^{2^{\alpha_k+1}-1}\frac{\left\|S_{n,n}f\right\|_{weak-L_p}^p\Phi\left(n\right)}{n^{3-2p}} \\
&\geq c_p\Phi\left(2^{\alpha_k}\right)\sum_{n=2^{\alpha_k-1}+1}^{2^{\alpha_k}-1}\frac{\left\|S_{2n+1,2n+1}f\right\|_{weak-L_p}^p}{\left(2n+1\right)^{3-2p}} \\
&\geq c_p\Phi\left(2^{\alpha_k}\right)\frac{2^{2\alpha_k(1-p)}}{\Phi^{1/4}\left(2^{\alpha_k}\right)}\sum_{n=2^{\alpha_k-1}+1}^{2^{\alpha_k}-1}\frac{1}{\left(2n+1\right)^{3-2p}} \\
&\geq c_p\Phi^{3/4}\left(2^{\alpha_k}\right)\to\infty, \qquad \text{when } k\to\infty.
\end{aligned}
$$

Combining (12-19) we complete the proof of Theorem 2.

References

[1] *G. Gát,* Inverstigations of certain operators with respect to the Vilenkin sistem, Acta Math. Hung., 61 (1993), 131-149.

[2] *G. Gát, U. Goginava, K. Nagy.* On the Marcinkiewicz-Fejér means of double Fourier series with respect to the Walsh-Kaczmarz system. Studia Sci. Math. Hungar. 46 (2009), no. 3, 399-421.

[3] *G. Gát, U. Goginava, G. Tkebuchava.* Convergence in measure of logarithmic means of quadratical partial sums of double Walsh-Fourier series. J. Math. Anal. Appl. 323 (2006), no. 1, 535-549.
[4] *U. Goginava,*The weak type inequality for the maximal operator of the Marcinkiewicz-Fejer means of the two-dimensional Walsh-Fourier series. J. Approximation Theory , 154, 2 (2008), 161-180.
[5] *U. Goginava, L. D. Gogoladze,* Strong Convergence of Cubic Partial Sums of Two-Dimensional Walsh-Fourier series, Constructive Theory of Functions, Sozopol 2010: In memory of Borislav Bojanov. Prof. Marin Drinov Academic Publishing House, Sofia, 2012, pp. 108-117.
[6] *L. D. Gogoladze,* On the strong summability of Fourier series, Bull of Acad. Scie. Georgian SSR, 52, 2 (1968), 287-292.
[7] *B. Golubov, A. Efimov* and *V. Skvortsov,* Walsh series and transformations, Dordrecht, Boston, London, 1991. Kluwer Acad. publ; 1991.
[8] *F. Schipp,* W.R. Wade, P. Simon and J. Pál, Walsh Series, an Introduction to Dyadic Harmonic Analysis. Adam Hilger, Bristol, New York, 1990.
[9] *P. Simon,* Strong convergence of certain means with respect to the Walsh-Fourier series, Acta Math. Hung. 49 (1987), 425-431.
[10] *P. Simon.* Strong Convergence Theorem for Vilenkin-Fourier Series. Journal of Mathematical Analysis and Applications, 245, (2000), pp. 52-68 .
[11] *B. Smith,* A strong convergence theorem for $H_1(T)$, in Lecture Notes in Math., 995, Springer, Berlin, (1994), 169-173.
[12] *G. Tephnadze,* On the Vilenkin-Fourier coefficients, Georgian Mathematical Journal, (to appear).
[13] *G. Tephnadze,* A note of the Fourier coefficients and partial sums of Vilenkin-Fourier series, Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis (AMAPN), (to appear).
[14] *F. Weisz,* Martingale Hardy spaces and their Applications in Fourier analysis, Springer, Berlin - Heidelberg - New York, 1994.
[15] *F. Weisz,* Summability of multi-dimensional Fourier series and Hardy space, Kluwer Academic, Dordrecht, Boston, London, 2002.
[16] *F. Weisz,* Strong convergence theorems for two-parameter Walsh-Fourier and trigonometric-Fourier series. (English) Stud. Math. 117, No.2, (1996), 173-194.

G. Tephnadze, Department of Mathematics, Faculty of Exact and Natural Sciences, Tbilisi State University, Chavchavadze str. 1, Tbilisi 0128, Georgia

E-mail address: giorgitephnadze@gmail.com